OCEAN**RAINFOREST**

VOLUME

☐ 1 gal / 3.78L
☐ 2.5 gal / 9.46 L
☐ ___ gal/ ___ L
8.43lb/gal @ 68˚F

SAKARI **BOOST**

CONTAINS
BENEFICIAL
SUBSTANCE
Kelp extract 99.9%



A PLANT BIOSTIMULANT derived from *Saccharina latissima* kelp intended to support a plant's natural nutrition process

REGISTERED ORGANIC INPUT MATERIAL cdfa

OMRI LISTED®

Guaranteed by:
Ocean Rainforest, Inc.
6710 Navigator Way
Goleta, California 93117 USA

STOP AND READ
THE LABEL BEFORE USING THE PRODUCT

LEARN MORE AT
oceanrainforest.com

Lot No.:
Manufacture Date:
Expiration Date: 2 YEARS AFTER THE MANUFACTURE DATE

PRECAUTIONS AND WARNINGS

- Do not eat, drink, or smokewhile using thisproduct
- Do not transport or store next to food products
- Keep out of reach of children
- Wash hands after using the product and before eating
- Do not ingest and avoid contact with eyes and skin

FIRST AID

If there is accidental skin or eye contact with this product, wash or rinse with water. If ingested, drink abundant water. In case of a more severe reaction, see a doctor and bring the product label.

STORAGE AND TRANSPORT

Theproductmustbe transported andstoredin its original container. Store the product in a ventilated, dry area that is protected from direct sunlight.

INSTRUCTIONS FOR USE

SakariBoost is a plant biostimulantmade from the seaweed *Saccharina latissima*, recommended for foliar application in combination with drench applications during the vegetative development of the crops listed in the table.
SakariBoost is compatible with most fungicides, insecticides, bactericides, and fertilizers. In case of doubt, a "jar" compatibility test is suggested.

WARRANTY NOTICE:

Asthe use of theproduct extendsbeyond the control of OCEAN RAINFOREST, INC., the company is not liable for misuse of the product or for any damages that may occur if the recommendations stated on this label are not followed.

DOSAGE RECOMMENDATIONS

*Additional applications should be made immediately prior to or following stress events such as heat, cold, or drought

CROP	APPLICATION	DOSAGE
FRUIT		
Avocado	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	70 fl oz/Ac
Berries	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	60 fl oz/Ac
Strawberries	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	60 fl oz/Ac
Citrus	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	70 fl oz/Ac
Grapes (wine, table, raisin, juice)	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	70 fl oz/Ac
Stone fruit	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	70 fl oz/Ac
VEGETABLES		
Brassica (cole) leafy vegetables, Bulb vegetables, Cucurbic vegetables, Fruiting vegetables	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	60 fl oz/Ac
NUTS		
Almonds	Soil application: Apply the product every two weeks to support nutrient uptake efficiency. Use the product before and during stress periods.	70 fl oz/Ac

Contact your local dealer or advisor for specific use recommendations.

OCEAN
RAINFOREST